

Free Energy of Gravitating Fermions

Neven Bilić¹ and Raoul D. Viollier²

¹Rudjer Bošković Institute, 10000 Zagreb, Croatia
bilic@thphys.irb.hr ; fax: 385-1-428 541

²Department of Physics, University of Cape Town
Rondebosch 7700, South Africa

February 7, 2008

Abstract

We study a system of self-gravitating massive fermions in the framework of the general-relativistic Thomas-Fermi model. We postulate the free energy functional and show that its extremization is equivalent to solving the Einstein's field equations. A self-gravitating fermion gas we then describe by a set of Thomas-Fermi type self-consistency equations.

Thermodynamical properties of the self-gravitating fermion gas have been extensively studied in the framework of the Thomas-Fermi model [1-6]. The system was investigated in the nonrelativistic Newtonian limit. The canonical and grand-canonical ensembles for such a system have been shown to have a nontrivial thermodynamical limit [1, 2]. Under certain conditions this system will undergo a phase transition that is accompanied by a gravitational collapse [2, 5] which may have important astrophysical and cosmological implications [6].

In this paper we formulate the general-relativistic version of the model. The effects of general relativity become important if the total rest-mass of the system is close to the Oppenheimer-Volkoff limit [7]. There are three main features that distinguish the relativistic Thomas-Fermi theory from the Newtonian one: *i*) the equation of state is relativistic *ii*) the temperature and chemical potential are metric dependent local quantities *iii*) the gravitational potential satisfies Einstein's field equations (instead of Poisson's equation).

Let us first discuss the general properties of a canonical, self-gravitating relativistic fluid. Consider a nonrotating fluid consisting of N particles in a spherical volume of radius R in

equilibrium at non-zero temperature. We denote by u_μ , p , ρ , n and σ the velocity, pressure, energy density, particle number density and entropy density of the fluid. Following Gibbons and Hawking [8] we postulate quite generally the free energy of the canonical ensemble as

$$F = M - \int_{\Sigma} T \sigma k^\mu d\Sigma_\mu, \quad (1)$$

where M is the total mass as measured from infinity, Σ is a spacelike hypersurface that contains the fluid and is orthogonal to the time-translation Killing vector field k^μ which is related to the velocity of the fluid

$$k^\mu = \xi u^\mu; \quad \xi = (k^\mu k_\mu)^{1/2}. \quad (2)$$

The entropy density of a relativistic fluid may be expressed as

$$\sigma = \frac{1}{T}(p + \rho - \mu n), \quad (3)$$

and T and μ are the local temperature and chemical potential. The metric generated by the mass distribution is static, spherically symmetric and asymptotically flat, i.e.

$$ds^2 = \xi^2 dt^2 - \lambda^2 dr^2 - r^2(d\theta^2 + \sin\theta d\phi^2). \quad (4)$$

ξ and λ may be represented in terms of the gravitational potential and mass

$$\xi = e^{\varphi(r)}, \quad (5)$$

$$\lambda = \left(1 - \frac{2m(r)}{r}\right)^{-1/2} \quad (6)$$

with

$$m(r) = \int_0^r dr' 4\pi r'^2 \rho(r'). \quad (7)$$

The equation of hydrostatic equilibrium [9, 10]

$$\partial_\nu p = -(p + \rho)\xi^{-1}\partial_\nu \xi, \quad (8)$$

together with the thermodynamic identity

$$d\frac{\mu}{T} = \frac{1}{n}d\frac{p}{T} + \frac{\rho}{n}d\frac{1}{T}, \quad (9)$$

implies

$$T\xi = T_0; \quad \mu\xi = \mu_0, \quad (10)$$

where T_0 and μ_0 are constants equal to the temperature and chemical potential at infinity. T_0 may be chosen arbitrarily, but μ_0 is an implicit functional of ξ due to the constraint that the total number of particles

$$\int_{\Sigma} n u^{\mu} d\Sigma_{\mu} = N \quad (11)$$

is fixed.

Based on Eq. (10) the free energy may be written in the form analogous to ordinary thermodynamics

$$F = M - T_0 S \quad (12)$$

with $M = m(R)$ and the total entropy S defined as

$$S = \int_0^R dr 4\pi r^2 \lambda \frac{1}{T} (p + \rho) - \frac{\mu_0}{T_0} N, \quad (13)$$

where we have employed the spherical symmetry to replace the proper volume integral as

$$\int_{\Sigma} u^{\mu} d\Sigma_{\mu} = \int_0^R dr 4\pi r^2 \lambda. \quad (14)$$

The following theorem shows that the quantity defined by Eq. (1) is indeed the free energy of a self-gravitating relativistic fluid at finite temperature.

Theorem 1 *Among all momentarily static, spherically symmetric configurations $\{\xi(r), m(r)\}$ which for given temperature T_0 at infinity contain a specified number of particles*

$$\int_0^R 4\pi r^2 dr \lambda(r) n(r) = N \quad (15)$$

within a spherical volume of given radius R , those and only those configurations that extremize the quantity F defined by (12) will satisfy the Einstein's field equation

$$\frac{d\xi}{dr} = \xi \frac{m + 4\pi r^3 p}{r(r - 2m)}, \quad (16)$$

with the boundary condition

$$\xi(R) = \left(1 - \frac{2M}{R}\right)^{1/2}. \quad (17)$$

Proof. From Eqs. (12) and (13) by making use of the identity (9), and the fact that $\delta(\mu/T) = \delta(\mu_0/T_0)$ and that N is fixed by the constraint (15), we find

$$\delta F = \delta M - \int_0^R dr 4\pi r^2 \frac{T_0}{T} (p + \rho) \delta \lambda - \int_0^R dr 4\pi r^2 \lambda \frac{T_0}{T} \delta \rho. \quad (18)$$

The variations $\delta \lambda$ and $\delta \rho$ can be expressed in terms of the variation $\delta m(r)$ and its derivative

$$\frac{d\delta m}{dr} = 4\pi r^2 \delta \rho. \quad (19)$$

This gives

$$\delta F = \delta M - \int_0^R dr 4\pi r^2 \frac{T_0}{T} (p + \rho) \frac{\partial \lambda}{\partial m} \delta m - \int_0^R dr \lambda \frac{T_0}{T} \frac{d\delta m}{dr}. \quad (20)$$

By partial integration of the last term and replacing T_0/T by ξ we find

$$\delta F = [1 - \lambda(R)\xi(R)] \delta M - \int_0^R dr \left[4\pi r^2 \xi (p + \rho) \frac{\partial \lambda}{\partial m} - \frac{d}{dr}(\lambda \xi) \right] \delta m, \quad (21)$$

where $\delta m(r)$ is an arbitrary variation on the interval $[0, R]$, except for the constraint $\delta m(0) = 0$. Therefore δF will vanish if and only if

$$4\pi r^2 \xi (p + \rho) \frac{\partial \lambda}{\partial m} - \frac{d}{dr}(\lambda \xi) = 0 \quad (22)$$

and

$$1 - \lambda(R)\xi(R) = 0. \quad (23)$$

Using Eqs. (6) and (7) we can write Eq. (22) in the form (16), and Eq. (23) gives the desired boundary condition (17). Thus, $\delta F = 0$ if and only if a configuration $\{\xi, m\}$ satisfies Eq. (16) with (17) as was to be shown.

Remark 1. A solutions to Eq. (16) is dynamically stable if the free energy assumes a minimum.

Remark 2. Our Theorem 1 is a finite temperature generalization of the result obtained for cold, catalyzed matter [11].

We now proceed to the formulation of the general-relativistic Thomas-Fermi model. Consider the case of a self-gravitating gas consisting of N fermions with the mass m_f contained in a sphere of radius R . Given the temperature at infinity T_0 , the following set of self-consistency equations defines the Thomas-Fermi equation:

1 Equation of state

$$n = g \int_0^\infty \frac{d^3 q}{(2\pi)^3} \frac{1}{1 + e^{E/T - \mu/T}}, \quad (24)$$

$$\rho = g \int_0^\infty \frac{d^3 q}{(2\pi)^3} \frac{E}{1 + e^{E/T - \mu/T}}, \quad (25)$$

$$p = gT \int_0^\infty \frac{d^3 q}{(2\pi)^3} \ln(1 + e^{-E/T + \mu/T}), \quad (26)$$

where g denotes the spin degeneracy factor, T and μ are local temperature and chemical potential, respectively, as defined in Eq. (10), and $E = \sqrt{m_f^2 + q^2}$.

2 Field equations

$$\frac{d\xi}{dr} = \xi \frac{m + 4\pi r^3 p}{r(r - 2m)}, \quad (27)$$

$$\frac{dm}{dr} = 4\pi r^2 \rho, \quad (28)$$

with the boundary conditions

$$\xi(R) = \left(1 - \frac{2m(R)}{R}\right)^{1/2}; \quad m(0) = 0. \quad (29)$$

3 Particle number constraint

$$\int_0^R dr 4\pi r^2 (1 - 2m/r)^{-1/2} n(r) = N. \quad (30)$$

One additional important requirement is that a solution of the self-consistency equations (24) to (30) have to minimize the free energy defined by Eq. (12)

Next we show that, in the Newtonian limit, we recover the usual Thomas-Fermi equation. Using the nonrelativistic chemical potential $\mu_{NR} = \mu_0 - m_f$ and the approximation $\xi = e^\varphi \simeq 1 + \varphi$, $E \simeq m_f + q^2/2m_f$ and $m/r \ll 1$, we find the usual Thomas-Fermi self-consistency equations [5, 6]

$$n = \frac{\rho}{m_f} = g \int_0^\infty \frac{d^3 q}{(2\pi)^3} \left(1 + \exp\left(\frac{q^2}{2m_f T_0} + \frac{m_f}{T_0} \varphi - \frac{\mu_{NR}}{T_0}\right)\right)^{-1}, \quad (31)$$

$$\frac{d\varphi}{dr} = \frac{m}{r^2}; \quad \frac{dm}{dr} = 4\pi r^2 \rho, \quad (32)$$

$$\varphi(R) = -\frac{m_f N}{R}; \quad m(0) = 0, \quad (33)$$

$$\int_0^R dr 4\pi r^2 n(r) = N. \quad (34)$$

The free energy (12) in the Newtonian limit yields

$$F = m_f N + \mu_{NR} N - \frac{1}{2} \int_0^R dr 4\pi r^2 n \varphi - \int_0^R dr 4\pi r^2 p \quad (35)$$

with

$$p = gT_0 \int_0^\infty \frac{d^3 q}{(2\pi)^3} \ln \left(1 + \exp\left(-\frac{q^2}{2m_f T_0} - \frac{m_f}{T_0} \varphi + \frac{\mu_{NR}}{T_0}\right) \right), \quad (36)$$

which, up to a constant, equals the Thomas-Fermi free energy [3].

A straightforward thermodynamic limit $N \rightarrow \infty$ as discussed by Hertel, Thirring and Narnhofer [2, 3] is in our case not directly applicable. First, in contrast to the non-relativistic case, there exist, at zero temperature, a limiting configuration with maximal M and N (Oppenheimer-Volkoff limit) and something similar can be expected at $T \neq 0$. Second, the scaling properties of the relativistic Thomas-Fermi equation are quite distinct from the non-relativistic one. The following scaling property can be easily shown:

If the configuration $\{\xi(r), m(r)\}$ is a solution to the self consistency equations (24) to (30), then the configuration $\{\xi = \xi(A^{-1}r), \tilde{m} = Am(A^{-1}r); A > 0\}$ is also a solution with the rescaled fermion number $\tilde{N} = A^{3/2}N$, radius $\tilde{R} = AR$, asymptotic temperature $\tilde{T}_0 = A^{-1/2}T_0$, and fermion mass $\tilde{m}_f = A^{-1/2}m_f$. The free energy is then rescaled as $\tilde{F} = AF$. Therefore, there exist a thermodynamic limit of $N^{-2/3}F$, with $N^{-2/3}R$, $N^{1/3}T_0$, $N^{1/3}m_f$ approaching constant values when $N \rightarrow \infty$. In that limit the Thomas-Fermi equation becomes exact. It is obvious that application of this model to stellar systems should work very well if the interactions among individual particles are negligible.

References

- [1] Thirring, W.: Z. Phys. **235** 339 (1970)
- [2] Hertel, P., Thirring, W.: Free Energy of Gravitating Fermions. Commun. Math. Phys. **24** 22-36 (1971); Hertel, P., Thirring, W. Thermodynamic Instability of a System of Gravitating Fermions. In Dürr, H.P. (ed.) Quanten und Felder. Braunschweig: Vieweg 1971
- [3] Hertel, P., Narnhofer, H. and Thirring, W.: Thermodynamic Functions for Fermions with Gravostatic and Electrostatic Interactions. Commun. Math. Phys. **28** 159-176 (1972)

- [4] Baumgartner, B.: Commun. Math. Phys. **48** 207-213 (1976)
- [5] Messer, J.: J. Math. Phys. **22** 2910-2917 (1981)
- [6] Bilić, N., Viollier, R.D.: Gravitational Phase Transition of Fermionic Matter. Cape Town preprint UCT-TP 229/96, astro-ph/9607077
- [7] Oppenheimer, J.R., Volkoff, G.M.: On Massive Neutron Cores Phys. Rev. **55** 374-381 (1939)
- [8] Gibbons, G.W., Hawking, S.W.: Action Integrals and Partition Function in Quantum Gravity Phys. Rev. **D55** 2752-2756 (1977)
- [9] Tolman, R.C.: Relativity Thermodynamics and Cosmology. Oxford: Clarendon Press 1934
- [10] Landau, L.D., Lifshitz, E.M.: Fluid Mechanics. Oxford: Pergamon Press 1959.
- [11] Harrison, B.K., Thorne, K.S., Wakano, M., Wheeler, J.A.: Gravitation Theory and Gravitational Collapse. Chicago: The University of Chicago Press 1965.